



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

09/869088

NL99/783

4

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98204371.3

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts:
im Auftrag


For the President of the European Patent Office

Le Président de l'Office européen des brevets
P.C.

Il.C. Hatten-Heckman
Il.C. Hatten-Heckman

HAAG,
HAGUE,
LA HAYE, LE

08/02/99





Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.
Application no.
Demande n°:

98204371.3

Anmeldetag:
Date of filing:
Date de dépôt:

22/12/98

Anmelder
Applicant(s)
Demandeur(s)
DSM N.V.

6411 TE Heerlen
NETHERLANDS

Bezeichnung der Erfindung
Title of the invention
Titre de l'invention

Process for the preparation of compounds with enhanced optical purity

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat
State
Pays

Tag
Date
Date

Aktenzeichen
File no.
Numéro de dépôt

Internationale Patentklassifikation
International Patent classification
Classification internationale des brevets

/

Am Anmeldetag benannte Vertragsstaaten
Contracting states designated at date of filing AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt


Bemerkungen
Remarks
Remarques



5

PROCESS FOR THE PREPARATION OF COMPOUNDS
WITH ENHANCED OPTICAL PURITY

10 The invention relates to a process for the preparation of a compound with enhanced optical purity wherein a mixture of the enantiomers of a chiral compound of formula 1:



20

wherein:

R_1 represents an alkyl or an aryl group

R_2 represents H, an alkyl or an aryl group

Y represents an alkyl group, an aryl group, $(CH_2)_nCOOH$,
 25 $(CH_2)_nCOOR$, $(CH_2)_nCONRR'$, CH_2OH , or $C\equiv N$ wherein R and
 R' independently represent H, an alkyl or aryl group,
 and n represents 0 or 1 is brought into contact with
 an enzyme having peptide deformylase activity with a
 bivalent metal ion as a cofactor wherein the metal is
 30 chosen from the groups 5-11 of the periodic system.

Enzymes having peptide deformylase activity
 are known in the literature e.g. from P. T. Ravi
 Rajagopalan et al., Biochemistry 1987, 26, 13910-13916
 wherein the use of peptide deformylase is described for
 35 the deformylation of several peptides with N
 formylmethionine at the N-terminus and of N
 formylmethionine. Although the known peptide
 deformylases showed reasonable deformylase activity
 when peptides were used as a substrate, they

showed no or little activity with respect to N-formylmethionine.

Applicant surprisingly found that the peptide deformylases having a bivalent metal ion as a cofactor according to the invention, did not only show a considerable activity towards the substrates according to formula 1, but also appeared to be enantioselective.

Substrates according to formula 1 that can be used in the process of the invention are for instance amino acids, for instance α - or β -amino acids with 1-20 C-atoms, in particular α -H- α -amino acids, α -methyl- α -amino acids, β -amino acids; esters of said amino acids wherein the ester group is for instance an alkyl group having 1-10 C-atoms; amides of said amino acids, wherein optionally the amide is N-substituted with 1 or 2, preferably 1, substituent chosen from alkyl or aryl, having 1-10 C-atoms; nitriles corresponding to said α -amino acids; amino alcohols corresponding to said α -amino acids; or amines for instance (optionally substituted) aromatic or aliphatic amines. Suitable substituents are for instance (optionally substituted) alkyl groups, for instance with 1-10 C-atoms.

In another embodiment of the present invention a mixture of the enantiomers of a (non protected) amino compound is subjected to a formylation in the presence of a peptide deformylase having a bivalent metal ion as a cofactor, wherein the metal is chosen from the groups 8-11 of the periodic system and a formylating agent, whereby one of the enantiomers is selectively converted into the corresponding N-formyl amino compound

Suitable formylating agents are for

instance formic acid in case a thermodynamically controlled formylation can be performed, or formic acid esters or amides when the formylation is kinetically controlled. In a thermodynamically controlled
5 formylation the equilibrium is shifted towards the side of formyl derivative, preferably by precipitation of the formyl derivative.

Peptide deformylases are in general enzymes having formyl methionine peptide deformylase activity.
10 It should be noticed that in the literature also other names are being used instead of the name Peptide deformylases; in particular the following names may be mentioned here: formylmethionine deformylase, N-formylmethionyl aminoacyl-tRNA deformylase, N-formyl-L-methionine amidohydrolase N-formylmethionyl-aminoacyl-tRNA amidohydrolase.
15

Suitable peptide deformylases to be used in the process according to the invention are peptide deformylases classified as EC 3.5.1.27. Preferably, the
20 enzyme is an enzyme having the activity as described for EC 3.5.1.27 because excellent results are being achieved in the deformylation with such enzymes. It should be noticed, that until recently it was believed that the enzyme coded as EC 3.5.1.31 is catalyzing a
25 different reaction. In the meantime however it has been shown that the enzymes known as EC 3.5.1.27 and EC 3.5.1.31 are coded for by exactly the same gene and have the same activity. Therefore, as used herein, the term EC 3.5.1.27 is encompassing not only EC 3.5.1.31
30 but likewise all other enzymes having the said activity, as described for EC 3.5.1.27.

Although the family of PDF's is composed of proteins with a relatively low level of sequence

identity. the 3D structures of the members of this family appear closely related one to each other with, in particular, the building of a common fold around the bivalent metal ion and three signature sequences. As is described (for PDF's indicated as PDF) by Wagner et al., J. Biol. Chem., 273, 11411-6 (1998), for many of these enzymes characteristically three short amino acid stretches are present as strictly conserved motifs, namely in that the enzymes contain the sequences (i) 10 HEXXH, (ii) EGCLS and (iii) GXGXAAXQ. In these sequences X represents any natural amino acid, and standard one letter codes for amino acids are used: A = alanine, C = cysteine, E = glutamic acid, G = glycine, H = histidine, L = leucine, S = serine and Q = 15 glutamine.

Peptide deformylases are obtainable for instance from eubacteria for example *Escherichia coli*, *Bacillus subtilis*, *Clostridium acetobutylicum*, *Clostridium beyerinckii*, *Haemophilus influenzae* 20 *Thermotoga maritima*, *Thermus aquaticus*, *Thermus thermophilus*, *Calothrix PCC 7601*, *Bacillus stearothermophilus* or *Lactococcus lactis*. Preferably an enzyme of *Escherichia Coli* is used.

The peptide deformylases according to the 25 invention require a bivalent metal ion whereby the metal is chosen from the groups 5-11 of the periodic system (New IUPAC version, see Handbook of Chemistry and Physics 70th edition, CRC Press, 1989-1990, inner page of cover), as a cofactor. Preferably the metal is 30 chosen from the group of V, Cr, Fe, Ni, Mn, Co, Cu, Pd and Pt, in particular from the group of Fe, Ni, Mn and Co.

Preferably the amount of the bivalent metal ions should be about equivalent to the number of moles

of enzyme. Suitably the molar ratio between these bivalent metal ions and the number of PDF molecules is in the range of 0.6 to 1.4, preferably of 0.8 to 1.2, and most preferred the amount of bivalent metal ions is
5 equimolar to the enzyme.

Exchange of the bivalent metal ions in the PDF's in order to obtain PDF enzymes with a co-factor as necessary for the present invention can be done by the various methods as described in Groche et al.,
10 Biochem. Biophys. Res. Comm., 246, 342-346, (1998). These methods include simple metal displacement by incubation of the native enzyme in an excess of the desired bivalent metal ion, if necessary preceded by the preparation of the apoenzyme via treatment of the
15 native enzyme with a metal chelation compound. Furthermore, the desired bivalent metal ion can already be introduced in (at least part of the enzyme molecules) by using a bacterial growth medium with an enhanced ratio of the desired bivalent metal ion over
20 Fe^{2+} .

In addition measures may be taken in order to enhance the stability of the enzyme, for instance the addition of stabilisation agents, for instance catalase, tris-(2-carboxyethyl)phosphine, glucose
25 oxidase, or combinations thereof; or enlarging the concentration of the PDF, for instance to a PDF concentration of at least 0.1 mg of PDF per ml, more preferably of least 1.0 mg/ml. The upper limit of the concentration of PDF is not critical if practical
30 concentrations are being used. The use of stabilisation measures is especially preferred when an easily

oxidisable metal ion, e.g. Fe^{2+} is present as a cofactor or an easily oxidisable substrate. If not, for instance in case Ni^{2+} is present as a cofactor the addition of a stabilisation agent appeared to be superfluous, as the
5 enzyme turned out to be very stable even without stabilisation agent.

The enzymes applied in the process according to the invention may be purified enzymes, a crude enzyme solution, microbial cells exhibiting the
10 required activity, a homogenate of cells or permeabilized cells. If required, the enzyme may be applied in an immobilized state or in a chemically modified form to ensure a good stability, reactivity and enantioselectivity of the enzymes under the
15 conditions utilized.

Alternatively, genetically engineered mutants of PDF's may be used which have for instance enhanced activity or enantioselectivity in the (de)formylation reaction. These mutants can be generated by a number of
20 different approaches; for instance, by site-directed mutagenesis, site-specific random mutagenesis, region-specific random mutagenesis, and completely random mutagenesis; the latter form of mutagenesis is better known as directed evolution. General applicable methods
25 to perform these different protein engineering approaches are well known to the skilled man. If a random approach will be applied, the mutagenesis cycle will need to be followed by selection of resistant and active mutant(s), thereby leading to the identification
30 of suitable mutants. To obtain PDF mutants also a combination of different protein engineering approaches and/or several rounds of random mutagenesis may be

used.

The reaction conditions for the enzymatic deformylation according to the invention are not very critical and may depend on the substrate used. Any suitable solvent system which is inert towards the PDF may be applied; such solvents include aqueous systems (solutions or slurries) or aqueous systems also containing a water-miscible organic solvent which is inert under the reaction conditions. Aqueous systems, however, are preferred. Also the concentration of the N-formyl compound is not critical, and may be for instance in the range of about 0.1 to 1000 mM. It is not necessary that all of the N-formyl compound is dissolved; part of it may be present as a slurry. The concentration of the PDF likewise is not very critical, and usually will be at 0.001 to 100 % by weight of the formyl compound, e.g. at about 0.2 mM of PDF. The pH for the reaction preferably is chosen in the range of 4.0 to 11.0, more preferably of 5.0 to 10.0. The temperature is not very critical, and suitably will be in the range of 10 to 50°C, e.g. at about 37°C, but for thermostable PDF enzymes higher temperatures may be applied.

In those cases wherein the absolute configuration of the (de)formylated enantiomer was determined, it appeared that the S-enantiomer was (de)formylated more rapidly than the R-enantiomer. The optical purity is given by the enantiomeric excess (ee), the enantioselectivity of the enzyme is represented by E, and calculated as k_2/k_1 , wherein k_2 is defined as the rate of (de)formylation of the most

rapidly (de)formylated enantiomer and k_1 is defined as the rate of (de)formylation of the least rapidly (de)formylated enantiomer.

Optionally a salt promoting hydrophobic interactions is added to the reaction mixture, for instance a sulphate, phosphate, sulphite or acetate of ammonium, Rb, K, Na, Cs or Li. Most preferably ammonium sulphate or lithium sulphate is used.

The invention will further be elucidated by the following examples, without being limited thereto.

Abbreviations:

TB medium: 12 g/l of Bacto-Tryptone, Difco; 24 g/l of yeast extract, Difco; 4 g/l of glycerole; 2.3 g/l of KH_2PO_4 ; 12.5 g/l of K_2HPO_4 ;

Hepes: N-2-hydroxyethylpiperazine-N'-2-ethane sulphuric acid;

AEBSF: 2-aminoethyl-p-benzene sulphonyl fluoride;

TCEP: tris-(2-carboxyethyl)-phosphine.

MOPS: 3-(N-morpholino)propane sulphonic acid

MES: 2-(N-morpholino)ethane sulphonic acid

Examples 1-15. Comparative experiments A and B

Isolation of PDF(Fe)

For a detailed discussion of the methods used reference is made to Groche et al., BBRC 246, 342-346 (1998).

PDF(Fe) was isolated from overproducing *E.coli* cells grown at 30°C in 1.6 l TB medium for 14-16

h. About 13 g (wet weight) cell paste were suspended in 26 ml buffer (20 mM Hepes/KOH, 100 mM KF, pH 7.7 supplemented with 10 µg/ml catalase from bovine liver (Boehringer Mannheim) and 1 mM AEBSF, disintegrated by sonication (Branson B12, 20 min) at 0°C and centrifuged at 200.000 g for 1 h. The clear supernatant (1.3 g of protein; according to biurete reaction) was mixed with 1.3 ml 10%(w/v) Polymix G-35 (BASF) adjusted to pH 7.7 and centrifuged at 40.000 g for 10 min. The supernatant was applied to a 20 ml Met-Lys-Sepharose column that had been equilibrated with 20 mM Hepes/KOH, 100 mM KF, 0.2 mM TCEP, pH 7.7. After washing with 120 ml of 20 mM Hepes/KOH, 100 mM KF, 0.2 mM TCEP, pH 7.7, PDF(Fe) was eluted with 150 ml 20 mM Hepes/KOH, 100 mM KCl, 0.2 mM TCEP, pH 7.7. The protein containing fractions were concentrated by ultrafiltration using an Amicon PM10 membrane (yield: 140 mg protein, 1400 U/mg; determined according to Groche et al.). After adjustment of the TCEP concentration to 1 mM and protein concentration to 40 mg/ml, the PDF(Fe) stock solution (40 mg/ml = 2 mM) was stored frozen at -60°C. After thawing, the PDF(Fe) stock solution could be used directly in the deformylation experiments described below. If however solutions with lower PDF(Fe) concentrations were required for these deformylation experiments, the PDF stock solution was diluted further in 20 mM Hepes/KOH, pH 7.7, 100 mM KCl, 1 mg/ml bovine serum albumin, 10 µg/ml catalase solution.

30 HPLC-analysis

In all cases HPLC conditions had to be

developed in which the two deformylated isomers were separated from each other and from the formylated isomers. To this end two different techniques were applied that is method 1 and method 2, as described
5 below.

From the quantities of deformylated isomers in the samples after various reaction times, both the initial deformylation rate constant (k_1 and k_2 in $M^{-1}s^{-1}$) could be calculated for both enantiomers, as well as
10 the respective ee values. The enantioselectivity of the enzyme (E value) was calculated by taking the ratio of k_1/k_2 and is given for all Examples in table 1, as well as the maximum ee value (ee_{max}) observed during the experiments.

15

Method 1 (without derivatization)

A Crownpak CR(+) column (4x150 mm) was used. Samples (5 μ l) withdrawn from the deformylation mixture were mixed with 95 μ l aqueous $HClO_4$ (10 mM) to
20 inactivate PDF(Fe). Following a brief centrifugation, 20 μ l of the supernatant were applied to the Crownpak CR(+) column. For specific chromatographic conditions and retention times see table 2.

25 Method 2 (Precolumn derivatization with o-phthalaldehyde (OPA) and N-acetyl-L-cysteine (NAC)). Samples (25 μ l) withdrawn from the deformylation mixture were mixed with 25 μ l aqueous $HClO_4$ (100 mM) to inactivate PDF(Fe). Following a brief centrifugation,
30 40 μ l of the supernatant were added to 80 μ l 1 M

aqueous $H_3BO_3/NaOH$ pH 9.4, subsequently 20 μl OPA reagent (consisting of OPA in H_2O/CH_3OH 1:1 v/v with a concentration as indicated in table 3) and 20 μl NAC reagent (consisting of NAC in H_2O/CH_3OH 1:1 v/v with a concentration as indicated in table 3) was added. After the time indicated in table 3 derivatization was terminated by addition of 80 μl (250 mM) aqueous H_3PO_4 , and 20 μl of the solution were instantaneously applied to a Nucleosil 120-5 C_{18} (250x4 mm) column. Temperature was always ambient and detection was spectrophotometric using a wavelength of 257 nm and/or 340 nm. The used eluent was a mixture of aqueous 0.05 M H_3PO_4 brought at pH 7.0 with 1 M NaOH, and a percentage of acetonitrile as indicated in table 3.

For derivatization of valine aminonitrile borate buffer was adjusted to pH 11 and addition of NAC reagent was done 10 min after OPA reagent had been added. Concentration of H_3PO_4 used for termination was 500 mM. Derivatization and separation conditions as well as the observed retention times for the deformedylated compounds analyzed are compiled in table 3.

Examples 1-12 and comparative experiments A and B were executed according to the procedures A, B, or C as given below as indicated in table 1. The results of the examples and comparative experiments are summarized in table 1 and the corresponding HPLC conditions in tables 2 and 3.

Method A

Deformylation in the presence of Li_2SO_4 at pH 7.2

Deformylation reactions were performed in 1.5 ml Eppendorf reaction test tubes. The reaction mixture with a total volume of 200 μ l contained 100 mM aqueous MOPS/NaOH, 2 M Li_2SO_4 buffer pH 7.2, and the concentration of formylated compound as indicated in table 1. After thermal equilibration to 37°C the deformylation reaction was started by the addition of the concentration of PDF as indicated in table 1. At various reaction times samples of the reaction mixture were withdrawn in which the reaction was stopped by addition of HClO_4 .

Method B

Deformylation in the absence of Li_2SO_4 at pH 7.2

Reactions were performed as described in Method A with the exception that 100 mM aqueous MOPS/NaOH, 250 mM NaCl, 0.1 mg/ml catalase buffer pH 7.2 was used in stead of 100 mM aqueous MOPS/NaOH, 2 M Li_2SO_4 buffer pH 7.2.

Method C

Deformylation in the absence of Li_2SO_4 at pH 6.2

Reactions were performed as described in Method A with the exception that 100 mM aqueous MES/NaOH buffer pH 6.2 was used in stead of 100 mM MOPS/NaOH, 2 M Li_2SO_4 buffer pH 7.2.

Table 1
Results of deformylation experiments

Ex.	Compound	Type of compound	Method	[Compound] (mM)	[PDP] (μ M)	k_2 ($M^{-1}s^{-1}$)	k_t ($M^{-1}s^{-1}$)	K	ee_{max} (%)
1	N-formyl-phenylglycine	α -H-amino acid	B	10	200	0.0047	10,6	2255	99,6
2	N-formyl-3-amino-3-phenyl-propionic acid	β -H-amino acid	B	10	10	<0.004	7,1	>1775	100
3	N-formyl-phenylglycine amide	α -H-amino acid amide	B	10	5.2	0.09	227	2522	99,7
4	N-formyl-tert-leucine amide	α -H-amino acid amide	A	4.8	200	0.0005	0,15	300	100
5	N-formyl- α -methyl-phenylglycine amide	α -alkyl-amino acid amide	A	10	200	0.0005	0,045	90	100
6	N-formyl-phenylglycinol	β -amino alcohol	B	10	200	0.029	0,69	23,8	90,5
7	N-formyl-phenylglycinol	β -amino alcohol	A	10	200	0.34	6,3	18,5	93,3
8	N-formyl-alaninol	β -amino alcohol	A	10	200	0.018	0,22	12	85,6
9	N-formyl-phenylalanine aminonitrile	α -aminonitrile	C	7.5	20	1	880	880	98,8
10	N-formyl-valine aminonitrile	α -aminonitrile	A	10	50	0.62	29,7	47,9	95,5
11	N-formyl- α -methoxy-phenylalanine aminonitrile	α -aminonitrile	B	7.2	2.5	2	1370	685	99,0
12	N-formyl-1-(1-naphthyl)-ethylamine	amine	A	0.42	200	0.03	0,45	15	90
A	N-acetyl-phenylglycine amide	α -H-amino acid amide	A	10	10	<0.001	<0.001	-	-
B	N-formyl-proline	α -H-imino acid	A	10	200	<0.004	<0.004	-	-

Table 2,
Analytical conditions and retention times analysed according to method 1

Ex.	Compound	Solvent	flow rate (ml/min)	T (°C)	Detection (nm)	Retention time (min)	Amine	formyl- compound
1	N-formyl-phenylglycine	10 mM aq. HClO ₄	1.0	40	210	2.1 (D)	3.8 (L)	9.6
2	N formyl 3-amino 3-phenyl propionic acid	85% 100 mM aq. HClO ₄ /15% CH ₃ OH	0.7	5	210	23.7	26.7	31.4
3	N-formyl-phenylglycine amide	10 mM aq. HClO ₄	0.8	22	210	3.2 (D)	12.6 (L)	6.3
6/7	N formyl phenylglycinol	95% 10 mM aq. HClO ₄ /5% CH ₃ OH	0.8	5	210	4.8 (L)	5.7 (D)	10.4
9	N formyl-phenylalanine aminonitrile	90% 10mM aq. HClO ₄ /10% CH ₃ OH	0.8	5	210	11.8	15.1	28.1
11	N formyl 4-methoxy-phenylalanine aminonitrile	90% 10mM aq. HClO ₄ /10% CH ₃ OH	0.8	5	210	23.8	30.7	52.0
12	N-formyl 1-(1 naphthyl)-ethylamine	85% 10 mM aq. HClO ₄ /15% CH ₃ OH	1.0	40	210	26.5 (S)	31.2 (R)	73.5
A	N acetylphenylglycine amide	10 mM aq. HClO ₄	0.8	22	210	3.2 (D)	12.6 (L)	6.9
B	N formyl proline	100 mM aq. HClO ₄	0.4	5	200	3.8	3.8	5.9

EP98204371.3

Table 31
Analytical conditions and retention times analyzed according to method 2

Ex. Compound	OPA (mg/ml)	NAC (mg/ml)	Time (min.)	% CH ₃ CN	Retention Time (min.)
2 N-formyl-3-amino-3-phenyl propionic acid	4	4	10	15	19.7
4 N-formyl tert leucine amide	4	8	10	22.5	14.9 (D)
5 N-formyl-n-butyl phenylglycine amide	16	16	30	20	24.4
8 N-formyl alaninol	4	4	5	15	16.9 (L)
10 N-formyl-valine amide/nitrile	16	4	5	20	8.6 (L)

CLAIMS

1. Process for the preparation of a compound with enhanced optical purity wherein a mixture of the enantiomers of a chiral compound of formula 1



wherein:

R_1 represents an alkyl or an aryl group

R_2 represents H, an alkyl or an aryl group

Y represents an alkyl group, an aryl group,

$(CH_2)_nCOCH$, $(CH_2)_nCOOR$, $(CH_2)_nCONRR'$, CH_2OH , or

$C \equiv N$ wherein R and R' independently represent H,

an alkyl or aryl group, and n represents 0 or 1,

is brought into contact with an enzyme having peptide deformylase activity with a bivalent

metal ion as a cofactor wherein the metal is

chosen from the groups 5-11 of the periodic

system.

2. Process for the preparation of a compound with enhanced optical purity wherein a mixture of the enantiomers of a chiral compound of formula 2



wherein:

R_1 represents an alkyl or an aryl group

R_2 represents H, an alkyl or an aryl group

Y represents an alkyl group, an aryl group,
(CH₂)_nCOOH, (CH₂)_n-COOR, (CH₂)_n-CONRR', CH₂OH, or
C≡N wherein R and R' represent H, an alkyl or
aryl group, and n represents 0 or 1, is subjected
to a formylation reaction in the presence of an
enzyme having peptide deformylase activity with a
bivalent metal ion as a cofactor wherein the
metal is chosen from the groups 5-11 of the
periodic system, and a formylating agent, whereby
one of the enantiomers is selectively converted
in the corresponding N-formyl compound

3. Process according to claim 2 wherein formic acid,
a formic acid amide or a formic acid ester is
used as a formylating agent.

4. Process according to any one of claims 1-3,
wherein the peptide deformylase is chosen from
the class EC 3.5.2.27 or EC 3.5.1.31.

5. Process according to any of claims 1-4, wherein
the peptide deformylase contains the sequences of
(i) HEXXH, (ii) EGCLS and (iii) GXGXAAXQ.

6. Process according to any of claims 1-5, wherein
the peptide deformylase is obtainable from
Escherichia coli.

7. Process according to any of claims 1-6, wherein
the bivalent metal is chosen from the group of
Fe, Ni, Mn and Co.

8. Process according to claim 7, wherein the
bivalent metal is Ni.


9. Process according to any of claims 1-8, wherein
in addition a stabilisation agent is added.

10. Process according to claim 9 wherein the
stabilisation agent is catalase.

11. Process according to claim 9 or 10 wherein the
bivalent metal is Fe.

ABSTRACT

5 Process for the preparation of a compound with enhanced optical purity wherein a mixture of the enantiomers of a chiral compound of formula :



wherein:

R_1 represents an alkyl or an aryl group

R_2 represents H, an alkyl or an aryl group

Y represents an alkyl group, an aryl group, $(CH_2)_nCOOH$,
 20 $(CH_2)_n-CCOR$, $(CH_2)_n-CONRR'$, CH_2CH , or $C \equiv N$ wherein R and R' independently represent H, an alkyl or aryl group, and n represents 3 or 1, is brought into contact with an enzyme having peptide deformylase activity with a bivalent metal ion as a cofactor wherein the metal is
 25 chosen from the groups 5-11 of the periodic system, or for the preparation of a formylated compound with enhanced optical purity from a mixture of the enantiomers of the corresponding not formylated chiral compound in the presence of a formylation agent.

30 Preferably the peptide deformylase is chosen from the class EC 3.5.2.27 or EC 3.5.1.31, and contains the sequences of (i) HEXXH, (ii) EGCLS and (iii) GXGXAAHQ. The bivalent metal may be chosen from the group of Fe, Ni, Mn and Co, preferably Ni or Fe.